



| | CHICKENS | | | | | |
|---|-------------|--------------|-------------------------|-----------------|------------------------|------------------------|
| | Egg Birds | | | | Meat Birds | |
| | 0 – 8 Weeks | 8 – 20 Weeks | 20 Weeks Through Laying | During Molt | 0 – 6 Weeks (Broilers) | 6 – 8 Weeks (Broilers) |
| Home Fresh Starter AMP | 3.0 – 5.0 | 13 – 16 | | | 8.0 – 10.0 | 5.5 – 7.0 |
| Home Fresh Grow & Show | | 14 – 18 | | | | |
| Home Fresh Extra Egg Layer | | | 0.20 – 0.25/day | 0.20 – 0.25/day | | |
| Home Fresh Finish | | | | | | 5.25 – 6.75 |
| Home Fresh Multi-Flock Chick N Game Starter/Grower 22 | 3.0 – 5.0 | | | | 8.0 – 10.0 | 5.5 – 7.0 |
| Home Fresh Better Feather | | | 0.20 – 0.25/day | 0.20 – 0.25/day | | |
| Home Fresh Organic Starter Broiler | 3.0 – 5.0 | | | | 8.0 – 10.0 | 5.5 – 7.0 |
| Home Fresh Organic Grower | | 14 – 18 | | | | |
| Home Fresh Organic Layer | | | 75 – 110 | | | |

Feed should be offered as free choice, but periodically make birds clean up. Ranges shown are due to the great variability in body size of breeds involved in each category.

For breeding flocks, restrict feed consumption to about 80% of normal consumption to prevent excess fattening. Provide clean, fresh water at all times.

*Feeding rates shown in above chart are approximate feed consumption in lb/bird during the total time frame in weeks as listed or as noted otherwise.